

ATLAS Trigger Status and Results From Commissioning Operations

Biglietti M.,^{68,38} Abolins M.,⁵⁶ Achenbach R.,³¹ Adorisio C.,¹⁵ Adragna P.,⁷⁶ Aharrouche M.,⁵⁷ Aielli G.,^{81,42} Al-Shabibi A.,³¹ Aleksandrov I.,²⁴ Alexandre G.,²⁶ Aloisio A.,^{68,38} Alviggi M.G.,^{68,38} Amorim A.,⁴⁹ Amram N.,⁸⁸ Anderson K.,¹⁶ Andrei V.,³¹ Anduaga X.,⁵⁰ Antonelli S.,^{10,36} Anulli F.,^{80,41} Aracena I.,⁸² Ask S.,⁵⁸ Asman B.,⁸⁶ Avolio G.,⁹¹ Backlund S.,¹⁴ Badescu E.,¹¹ Baines J.,⁷⁷ Banerjee S.,⁹⁸ Barnett B.M.,⁷⁷ Bartoldus R.,⁸² Batraneanu S.,^{12,14} Battaglia A.,⁷ Bauss B.,⁵⁷ Beauchemin P.,⁷² Beck H.P.,⁷ Bee C.,⁵⁹ Behera P.,⁴⁵ Bell P.,⁵⁸ Bell W.H.,²⁸ Bellagamba L.,^{10,36} Bellomo M.,³⁹ Ben Ami S.,⁸⁷ Bendel M.,⁵⁷ Benhammou Y.,⁸⁸ Benslama K.,⁷⁸ Berge D.,¹⁴ Berger N.,⁴⁸ Bernius C.,⁹² Berry T.,³⁴ Bianco M.,^{52,37} Biau C.,⁹⁵ Bizjak I.,⁹² Blair R.E.,¹ Bogaerts A.,¹⁴ Bohm C.,⁸⁶ Boisvert V.,³⁴ Bold T.,^{91,20} Bondioli M.,⁹¹ Booth J.R.A.,⁸ Borer C.,⁷ Boscherini D.,^{10,36} Bosman M.,⁵ Boyd J.,¹⁴ Bracinik J.,⁸ Brawn I.P.,⁷⁷ Brelier B.,⁶² Bressler S.,⁸⁷ Brock R.,⁵⁶ Brooks W.K.,⁹⁷ Brunet S.,²¹ Bruni A.,^{10,36} Bruni G.,^{10,36} Buccif F.,²⁶ Buda S.,¹¹ Burckhart-Chromek D.,¹⁴ Buttar C.,²⁸ Camarri P.,^{81,42} Campanelli M.,⁹² Canale V.,^{68,38} Capasso L.,^{68,38} Caprini M.,¹¹ Caracinha D.,⁵³ Caramarcu C.,¹¹ Cardarelli R.,^{81,42} Carlino G.,³⁸ Casadei D.,⁶⁵ Casado P.,⁵ Cataldi G.,³⁷ Cerri A.,¹⁴ Charlton D.G.,⁸ Childers J.T.,³¹ Chiodini G.,^{52,37} Ciapetti G.,^{80,41} Cimino D.,⁴⁰ Ciobotaru M.,^{91,12} Coccaro A.,²⁷ Collins N.J.,⁸ Conde Muino P.,⁴⁹ Conventi F.,³⁸ Corradi M.,^{10,36} Corso-Radu A.,⁹¹ Costa M.J.,⁹⁶ Coura Torres R.,⁷⁹ Cranmer K.,⁶⁵ Crone G.,⁹² Crupi R.,^{52,37} Cuenca Almenar C.,⁹¹ Curtis C.J.,⁸ Czyczula Z.,⁶⁴ Dam M.,⁶⁴ Damazio D.,⁴ Darlea G.L.,¹⁴ Davis A.O.,⁷⁷ De Pedis D.,^{80,41} De Santo A.,³⁴ de Seixas J.M.,⁷⁹ DeAsmundis R.,³⁸ Degenhardt J.,⁷⁵ della Volpe D.,^{68,38} Della Pietra M.,^{68,38} Delsart P.-A.,⁶² Demers S.,⁸² Demirkoz B.,¹⁴ Di Mattia A.,⁵⁶ Di Nardo R.,^{81,42} Diaz M.,⁸⁴ DiCiaccio A.,^{81,42} Dionisi C.,^{80,41} Djilkibaev R.,⁶⁵ Dobson E.,⁷² Dobson M.,¹⁴ Dogaru M.,¹¹ dos Anjos A.,¹⁰⁰ Dotti A.,⁴⁰ Dova M.T.,⁵⁰ Drake G.,¹ Dufour M.-A.,⁶⁰ Eckweiler S.,⁵⁷ Ehrenfeld W.,^{30,21} Eifert T.,²⁶ Eisenhandler E.,⁷⁶ Ellis N.,¹⁴ Emeliyanov D.,⁷⁷ Enoque Ferreira de Lima D.,⁷⁹ Ermoline Y.,⁵⁶ Etzion E.,⁸⁸ Falciano S.,⁴¹ Farthouat P.,¹⁴ Faulkner P.J.W.,⁸ Fedorko A.,¹⁴ Fedorko I.,¹⁴ Feng E.,¹⁶ Ferland J.,⁶² Ferrag S.,²⁸ Ferrari R.,⁷⁴ Ferrer M.L.,²⁵ Fischer G.,²¹ Flacher H.,¹⁴ Fleckner J.E.,⁵⁷ Flowerdew M.,⁵⁴ Föhlich F.,³¹ Fonseca-Martin T.,³⁴ Francis D.,¹⁴ Fratina S.,⁷⁵ Fukunaga C.,⁹⁰ Gadomski S.,²⁶ Gallacher M.P.,⁸ Garitaonandia Elejabarrieta H.,⁶⁹ Gaudio G.,⁷⁴ Gee C.N.P.,⁷⁷ George S.,³⁴ Ghete V.M.,⁴⁴ Giagu S.,^{80,41} Gillman A.R.,⁷⁷ Giusti P.,^{10,36} Goebel M.,²¹ Gonçalo R.,³⁴ Gonzalez Pinto F.,¹³ Gonzalez Silva L.,¹³ Göringer C.,⁵⁷ Gorini B.,¹⁴ Gorini E.,^{52,37} Grabowska-Bold I.,^{91,20} Grancagnolo S.,³⁵ Green B.,³⁴ Groll M.,⁵⁷ Guida A.,^{52,37} Haas S.,¹⁴ Hadavand H.,⁸³ Hadley D.R.,⁸ Haller J.,^{30,21} Hamilton A.,²⁶ Hanke P.,³¹ Hansen J.R.,⁶⁴ Hasagawa S.,⁶⁷ Hasegawa Y.,⁸⁵ Hauschild M.,¹⁴ Hauser R.,⁵⁶ Hayakawa T.,⁴⁷ Head S.,⁵⁸ Heijboer A.,¹⁴ Heim S.,⁵⁶ Hellman S.,⁸⁶ Hershenhorn A.,⁸⁷ Hidvégi A.,⁸⁶ Hillier S.J.,⁸ Hirayama S.,⁸⁹ Hod N.,⁸⁸ Hori T.,⁴⁷ Hryn'ova T.,⁴⁸ Hughes-Jones R.,⁵⁸ Huston J.,⁵⁶ Iacobucci G.,^{68,38} Idarraga J.,⁶² Iengo P.,^{68,38} Igonkina O.,⁶⁹ Ikeno M.,⁴⁶ Ishikawa A.,⁴⁷ Ishino M.,⁸⁹ Iwasaki H.,⁴⁶ Izzo V.,⁶⁸ Jain V.,⁴³ Jez P.,⁶⁴ Jimenez Otero S.,¹⁴ Johansen M.,⁸⁶ Johns K.,³ Jones G.,⁵⁸ Joos M.,¹⁴ Kadosaka T.,⁴⁷ Ka jomovitz E.,⁸⁷ Kalinowski A.,⁷⁸ Kama S.,²¹ Kanaya N.,⁸⁹ Kanega F.,⁸⁹ Kaushik V.,³ Kawagoe K.,⁴⁷ Kawamoto T.,⁸⁹ Kazarov A.,⁷³ Kehoe R.,⁸³ Kessoku K.,⁸⁹ Khorauli G.,⁷⁸ Kieft G.,⁶⁹ Kirk J.,⁷⁷ Kiyamura H.,⁴⁷ Klovfer P.,¹⁴ Klous S.,⁶⁹ Kluge E.-E.,³¹ Kobayashi T.,⁸⁹ Koeneke K.,²¹ Kolos S.,^{91,73} Kono T.,¹⁴ Konoplich R.,⁶⁵ Konstantinidis N.,⁹² Korcyl K.,¹⁹ Kordas K.,⁷ Kotov V.,²⁴ Kowalewski R.V.,⁹⁸ Krasznahorkay A.,^{14,22} Kreisel A.,⁸⁸ Kubota T.,⁸⁹ Kugel A.,³² Kurashige H.,⁴⁷ Kwee R.,³⁵ Landon M.,⁷⁶ Lane J.,⁵⁸ Lankford A.J.,⁹¹ Leahu L.,^{12,14} Leahu M.,¹² LeCompte T.,¹ Ledroit F.,²⁹ Lehmann Miotto G.,¹⁴ Lei X.,³ Lellouch D.,⁹⁹ Lendermann V.,³¹ Levinson L.,⁹⁹ Li S.,²¹ Liberti B.,^{81,42} Lilley J.N.,⁸ Lim H.,¹ Linnemann J.T.,⁵⁶ Lohse T.,³⁵ Losada M.,⁹ Lowe A.,⁴³ Luci C.,^{80,41} Luminari L.,⁴¹ Lundberg J.,¹⁴ Lupu N.,⁸⁷ Mackeprang R.,¹⁴ Maettig S.,^{30,21} Mahboubi K.,³¹ Mahout G.,⁸ Maltrana D.,⁸⁴ Mamuzic J.,²¹ Männer R.,³² Mapelli L.,¹⁴ Marchese F.,^{81,41} Marino C.,⁴³ Martin B.,¹⁴ Martin B.T.,⁵⁶ Martyniuk A.,⁵⁸ Marzano F.,⁴¹ Masik J.,⁵⁸ Matsushita T.,⁴⁷ McMahon T.,³⁴ Mcpherson R.,⁹⁸ Medinnis M.,²¹ Meessen C.,⁵⁹ Meier K.,³¹ Meirosu C.,¹² Merola L.,^{68,38} Messina A.,¹⁴ Middleton R.P.,⁷⁷ Mikenberg G.,⁹⁹ Mincer A.,⁶⁵ Mineev M.,²⁴ Misiejuk A.,³⁴

Moa T.,⁸⁶ Moenig K.,²¹ Monticelli F.,⁵⁰ Mora Herrera C.,²⁶ Moreno D.,⁹ Morettini P.,²⁷ Morris J.D.,⁷⁶
 Müller F.,³¹ Murillo Garcia R.,⁹¹ Nagano K.,⁴⁶ Nagasaka Y.,³³ Nakatsuka H.,⁴⁷ Navarro G.A.,⁹ Negri
 A.,⁷⁴,³⁹ Nemethy P.,⁶⁵ Neusiedl A.,⁵⁷ Nisati A.,⁴¹ Niwa T.,⁴⁷ Nomoto H.,⁸⁹ Nozaki M.,⁴⁶ Nozicka M.,²¹
 Nurse E.,⁹² Ochi A.,⁴⁷ Oda S.,⁸⁹ Ohm C.,⁸⁶ Okumura Y.,⁶⁷ Oltmann B.,⁵⁷ Olvito D.,⁷⁵ Omachi C.,⁴⁷
 Osculati B.,²⁷ Oshita H.,⁸⁵ Osuna C.,⁵ Owen M.A.,⁵⁸ Özcan E.,⁹² Ozone K.,⁴⁶ Padilla C.,⁵ Panes B.,⁸⁴
 Panikashvili N.,⁶¹ Parodi F.,²⁷ Pasqualucci E.,⁴¹ Pastore F.,¹⁴ Patricelli S.,⁶⁸,³⁸ Pauly T.,¹⁴ Perantoni M.,⁷⁹
 Perera V.J.O.,⁷⁷ Perez E.,⁵ Perez Reale V.,¹⁸ Perrino R.,⁵²,³⁷ Pessoa Lima Junior H.,⁷⁹ Petcu M.,¹¹ Petersen
 B.,¹⁴ Petersen J.,¹⁴ Petrolo E.,⁸⁰,⁴¹ Piegaia R.,¹³ Pilcher J.,¹⁶ Pinzon G.,⁹ Policicchio A.,¹⁵ Polini A.,¹⁰,³⁶
 Pope B.G.,⁵⁶ Potter C.,⁶⁰ Prieur D.P.F.,⁷⁷ Primavera M.,³⁷ Prokoshin F.,⁹⁷ Qian W.,⁷⁷ Quinonez F.,⁸⁴ Ra
 jagopalan S.,⁴ Ramos Dos Santos Neves R.,⁴⁹ Reinherz E.,⁸⁸ Reinsch A.,⁷⁰ Renkel P.,⁸³ Rescigno M.,⁴¹
 Rieke S.,⁵⁷ Risler C.,³⁵ Riu I.,⁵ Robertson S.,⁶⁰ Roda C.,⁴⁰ Rodriguez D.,⁹ Rogriquer Y.,⁹ Roich A.,⁹⁹
 Romeo G.,¹³ Rosati S.,⁸⁰,⁴¹ Rühr F.,³¹ Ruiz-Martinez A.,⁹⁶ Ryabov Y.,⁷³ Ryan P.,⁵⁶ Saavedra A.,⁹³
 Sakamoto H.,⁸⁹ Salamon A.,⁸¹,⁴² Salnikov A.,⁸² Salvatore D.,¹⁵ Salvatore F.,³⁴ Sankey D.P.C.,⁷⁷
 Santamarina C.,⁶⁰ Santonico R.,⁸¹,⁴² Sarkisyan-Grinbaum E.,⁹⁴ Sasaki O.,⁴⁶ Savu D.,¹⁴ Scannicchio D.,¹⁴
 Schäfer U.,⁵⁷ Schiavi C.,²⁷ Schlereth J.,¹ Schmitt K.,³¹ Schmitt S.,²¹ Scholtes I.,¹⁴ Schooltz D.,⁵⁶ Schroer
 N.,³² Schultz-Coulon H.-C.,³¹ Schwienhorst R.,⁵⁶ Scott W.,⁷⁷ Segura E.,⁵ Sekhniaidze G.,³⁸ Sherman D.,¹⁴
 Shimo jima M.,⁶⁶ Sidoti A.,³⁵ Silverstein D.,⁸² Silverstein S.,⁸⁶ Sinev N.,⁷⁰ Siragusa G.,⁵⁷ Sivoklov S.,⁶³
 Sjoen R.,⁶ Sjölin J.,⁸⁶ Sloper J.E.,¹⁴ Smizanska M.,⁵¹ Soffer A.,⁸⁸ Soloviev I.,⁷³ Spagnolo S.,⁵²,³⁷ Spiwoкс
 R.,¹⁴ Staley R.J.,⁸ Stamen R.,³¹ Stancu S.,⁹¹,¹² Steinberg P.,⁴ Stelzer J.,²¹ Stockton M.C.,⁸ Stradling A.,¹⁰⁰
 Straessner A.,²³ Strom D.,⁷⁰ Su D.,⁸² Sugaya Y.,⁷¹ Sugimoto T.,⁶⁷ Sushkov S.,⁵ Sutton M.,⁹² Suzuki Y.,⁴⁶
 Szymocha T.,¹⁹ Taffard A.,⁹¹ Takahashi Y.,⁶⁷ Takeda H.,⁴⁷ Takeshita T.,⁸⁵ Tamsett M.,³⁴ Tan C.L.A.,⁸
 Tanaka S.,⁴⁶ Tapprogge S.,⁵⁷ Tarem S.,⁸⁷ Tarem Z.,⁸⁷ Taylor C.,⁹² Taylor R.P.,⁹⁸ Teixeira-Dias P.,³⁴
 Thomas J.P.,⁸ Thompson P.D.,⁸ Tokushuku K.,⁴⁶ Tollefson K.,⁵⁶ Tomoto M.,⁶⁷ Topfel C.,⁷ Torrence E.,⁷⁰
 Touchard F.,⁵⁹ Tremblet L.,¹⁴ Tripiana M.,⁵⁰ Tsuno S.,⁴⁶ Ünel G.,⁹¹ Urquijo P.,²⁶ Urrejola P.,⁸⁴ Usai G.,¹⁶
 Vachon B.,⁶⁰ Vandelli W.,¹⁴ Vari R.,⁸⁰,⁴¹ Vaz Gil Lopes L.,¹⁷ Veneziano S.,⁸⁰,⁴¹ Ventura A.,⁵²,³⁷ Vercesi
 V.,⁷⁴ Vermeulen J.C.,⁶⁹,² Von Der Schmitt J.,⁵⁵ Vorwerk V.,⁵ Wang M.,⁷⁸ Watkins P.M.,⁸ Watson A.T.,⁸
 Weber P.,³¹ Weidberg T.,⁷² Wengler T.,⁵⁸ Werner P.,¹⁴ Werth M.,⁹¹ Wessels M.,³¹ Wheeler-Ellis S.,⁹¹
 Whiteson D.,⁹¹ Wickens F.J.,⁷⁷ Wiedenmann W.,¹⁰⁰ Wielers M.,⁷⁷ Wildt M.,⁵⁷ Wilkens H.,¹⁴ Winklmeier
 F.,¹⁴ Wu S.-L.,¹⁰⁰ Wu X.,²⁶ Xella S.,⁶⁴ Yamazaki Y.,⁴⁷ Yang U.,⁵⁸ Yasu Y.,⁴⁶ Zanello L.,⁸⁰,⁴¹ Zema F.,¹⁴
 Zhang J.,¹ Zhao L.,⁶⁵ Zobernig H.,¹⁰⁰ zur Nedden M.,³⁵

¹ Argonne National Laboratory, Argonne, Illinois

² University of Amsterdam, Amsterdam

³ University of Arizona, Tucson, Arizona

⁴ Brookhaven National Laboratory (BNL), Upton, New York

⁵ Institut de Física d'Altes Energies (IFAE), Universitat Autònoma de Barcelona, Bellaterra (Barcelona)

⁶ University of Bergen, Bergen

⁷ Laboratory for High Energy Physics, University of Bern, Bern

⁸ School of Physics and Astronomy, The University of Birmingham, Birmingham

⁹ Universidad Antonio Narino, Bogotá, Colòmbia

¹⁰ Università di Bologna, Dipartimento di Fisica, Bologna

¹¹ National Institute of Physics and Nuclear Engineering, Bucharest

¹² UPB - 'Politehnica' University of Bucharest, Bucharest

¹³ Universidad de Buenos Aires, Buenos Aires

¹⁴ European Laboratory for Particle Physics (CERN), Geneva

¹⁵ Dipartimento di Fisica dell' Università della Calabria e I.N.F.N., Cosenza

¹⁶ University of Chicago, Enrico Fermi Institute, Chicago, Illinois

¹⁷ Laboratório de Instrumentação e Física Experimental de Partículas, Coimbra

¹⁸ Nevis Laboratory, Columbia University, Irvington, New York

¹⁹ Institute of Nuclear Physics, Polish Academy of Sciences, Cracow

²⁰ Faculty of Physics and Applied Computer Science, AGH University of Science and Technology, Cracow

- ²¹ Deutsches Elektronen-Synchrotron (DESY), Hamburg and Zeuthen
²² University of Debrecen
²³ Technical University Dresden, Dresden
²⁴ Joint Institute for Nuclear Research, Dubna
²⁵ Laboratori Nazionali di Frascati dell' I.N.F.N., Frascati
²⁶ Section de Physique, Université de Genève, Geneva
²⁷ Dipartimento di Fisica dell' Università di Genova e I.N.F.N., Genova
²⁸ Department of Physics and Astronomy, University of Glasgow, Glasgow
²⁹ Laboratoire de Physique Subatomique et de Cosmologie de Grenoble (LPSC), IN²P³-CNRS- Université Joseph Fourier, Grenoble
³⁰ University of Hamburg, Germany
³¹ Kirchhoff Institut für Physik, Universität Heidelberg, Heidelberg
³² ZITI Ruprecht-Karls-University Heidelberg
³³ Hiroshima Institute of Technology, Hiroshima
³⁴ Department of Physics, Royal Holloway, University of London, Egham
³⁵ Institut für Physik, Humboldt-Universität zu Berlin
³⁶ I.N.F.N. Bologna
³⁷ I.N.F.N. Lecce
³⁸ I.N.F.N. Napoli
³⁹ I.N.F.N. Pavia
⁴⁰ Dipartimento di Fisica dell' Università di Pisa e I.N.F.N., Pisa
⁴¹ I.N.F.N. Roma
⁴² I.N.F.N. Tor Vergata
⁴³ Indiana University, Bloomington, Indiana
⁴⁴ Institute for Astro- and Particle Physics, Leopold-Franzens-Universität Innsbruck
⁴⁵ Iowa State University, Ames, Iowa
⁴⁶ KEK, High Energy Accelerator Research Organisation, Tsukuba
⁴⁷ Kobe University, Kobe
⁴⁸ Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), IN²P³-CNRS, Annecy-le-Vieux
⁴⁹ Laboratório de Instrumentação e Física Experimental, Lisboa
⁵⁰ Universidad Nacional de La Plata, La Plata
⁵¹ Department of Physics, Lancaster University, Lancaster
⁵² Università degli Studi del Salento, Dipartimento di Fisica, Lecce
⁵³ University Católica-Figueira da Foz and University Nova de Lisboa, Lisbon
⁵⁴ University of Liverpool, Liverpool
⁵⁵ Max-Planck-Institut für Physik, München
⁵⁶ Michigan State University, Department of Physics and Astronomy, East Lansing, Michigan
⁵⁷ Institut für Physik, Universität Mainz, Mainz
⁵⁸ School of Physics and Astronomy, University of Manchester, Manchester
⁵⁹ Centre de Physique des Particules de Marseille, IN²P³-CNRS, Marseille
⁶⁰ Department of Physics, McGill University, Montreal
⁶¹ University of Michigan, Department of Physics, Ann Arbor, Michigan
⁶² University of Montreal, Montreal
⁶³ Moscow State University, Moscow
⁶⁴ Niels Bohr Institute, University of Copenhagen, Copenhagen
⁶⁵ Department of Physics, New York University, New York
⁶⁶ Nagasaki Institute of Applied Science, Nagasaki
⁶⁷ Nagoya University, Nagoya
⁶⁸ Università di Napoli 'Federico II', Dipartimento di Scienze Fisiche, Napoli
⁶⁹ Nikhef National Institute for Subatomic Physics, Amsterdam
⁷⁰ University of Oregon, Eugene, Oregon
⁷¹ Osaka University, Osaka

- ⁷² Department of Physics, Oxford University, Oxford
- ⁷³ Petersburg Nuclear Physics Institute (PNPI), St. Petersburg
- ⁷⁴ Università di Pavia, Dipartimento di Fisica Nucleare e Teorica, Pavia
- ⁷⁵ Department of Physics, University of Pennsylvania, Philadelphia, Pennsylvania
- ⁷⁶ Physics Department, Queen Mary, University of London, London
- ⁷⁷ STFC Rutherford Appleton Laboratory, Harwell Science and Innovation Campus, Didcot, Oxon
- ⁷⁸ University of Regina, Regina
- ⁷⁹ Universidade Federal do Rio de Janeiro, COPPE/EE/IF, Rio de Janeiro
- ⁸⁰ Università di Roma I 'La Sapienza', Dipartimento di Fisica, Roma
- ⁸¹ Dipartimento di Fisica dell' Università di Roma II 'Tor Vergata'
- ⁸² Stanford Linear Accelerator Center (SLAC), Stanford
- ⁸³ Department of Physics, Southern Methodist University, Dallas, Texas
- ⁸⁴ Pontificia Universidad Católica, Santiago
- ⁸⁵ Faculty of Science, Shinshu University, Matsumoto
- ⁸⁶ Fysikum, Stockholm University, Stockholm
- ⁸⁷ Department of Physics, Technion, Haifa
- ⁸⁸ School of Physics and Astronomy, Tel Aviv University, Tel Aviv
- ⁸⁹ International Center for Elementary Particle Physics, University of Tokyo, Tokyo
- ⁹⁰ Physics Department, Tokyo Metropolitan University, Tokyo
- ⁹¹ Department of Physics and Astronomy, University of California, Irvine, California
- ⁹² Department of Physics and Astronomy, University College London, London
- ⁹³ University of Sydney, Sydney
- ⁹⁴ Department of Physics, The University of Texas at Arlington, Arlington, Texas
- ⁹⁵ Université de Technologie de Belfort-Montbéliard
- ⁹⁶ Instituto de Física Corpuscular (IFIC) Universidad de Valencia
- ⁹⁷ Universidad Técnica Federico Santa María, Valparaíso ⁹⁸ University of Victoria, Victoria
- ⁹⁹ Department of Particle Physics, The Weizmann Institute of Science, Rehovot
- ¹⁰⁰ Department of Physics, University of Wisconsin, Madison, Wisconsin
- ¹⁰¹ Università di Napoli 'Parthenope'

The ATLAS trigger system is designed to select rare physics processes of interest from an extremely high rate of proton-proton collisions, reducing the LHC incoming rate of about 10^7 . The short LHC bunch crossing period of 25 ns and the large background of soft-scattering events overlapped in each bunch crossing pose serious challenges, both on hardware and software, that the ATLAS trigger must overcome in order to efficiently select interesting events. The ATLAS trigger consists of hardware based Level-1, and a two-level software based High-Level Trigger (HLT). Data bandwidth and processing times in the higher level triggers are reduced by region of interest guidance in the HLT reconstruction steps. High flexibility is critical in order to adapt to the changing luminosity, backgrounds and physics goals. It is achieved by the use of inclusive trigger menus and modular software design. Selection algorithms have been developed which provide the required elasticity to detect different physics signatures and to control the trigger rates. In this paper an overview of the ATLAS trigger design, status and expected performance, as well as the results from the on-going commissioning with cosmic rays and first LHC beams, is presented.

XII Advanced Computing and Analysis Techniques in Physics Research
Erice, Italy
3-7 November, 2008

1. Introduction

The ATLAS experiment [1, 2] is one of the four experiments at the Large Hadron Collider (LHC), that is scheduled to start to operate in the year 2009. At present, the installation of the different components of the ATLAS detector is completed in the underground cavern and the commissioning process has already started. At the design luminosity of $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ the detector will be exposed every 25 ns to, on average, 25 proton-proton interactions. These conditions make the environment for the trigger system extremely demanding. It has to reduce the input event rate of 1 GHz to the limit of the storage rate of 200 Hz or less. The ATLAS trigger is composed of three levels: Level 1 (L1) [3] which is hardware-based, the Level 2 (L2) and Event Filter (EF) (collectively referred to as the High Level Trigger or HLT [4]) based on software algorithms analyzing the data on large computing farms. This note focuses the trigger strategy that can be deployed during the initial low luminosity running and describe the commissioning operations of the Trigger for initial running with cosmic rays and first LHC beams.

2. The ATLAS Detector and Trigger Strategy

In order to enrich the new-physics content of the recorded data (and remain within the limits of technology to data transfer) a data selection is made through the use of a highly-selective three-level trigger. L1 reduces the 40 MHz input rate (bunch-crossing rate) to about 75 kHz (upgradable to 100 kHz). It uses coarse granularity data from the muon and calorimeter detector systems. L1 must reach a decision within $2.2\mu\text{s}$. The LVL1 selection is mainly based on the identification of high transverse momentum objects in the detector.

The allowed L1 configurations (also called L1 items) are programmed in the Central Trigger Processor (CTP) [5]; each of L1 items is a logical combination of a given multiplicities of one or more of the configured L1 thresholds.

The L2 reconstruction is seeded by L1 Regions of Interest (RoIs) and has access to the full detector granularity. The size of the RoI is determined by the L2 algorithms depending on the type of triggered object. RoI mechanism is used to minimize the amount of data needed to calculate the trigger decisions thus reducing the overall network data traffic considerably. The trigger requests only a few percent of the detector data, leading to large savings in the necessary network bandwidth. Within each RoI, L2 reconstructs physics objects using fast algorithms that use detector information not available at L1, like Muon Spectrometer precision chambers and the Inner Detector. The information from individual sub-systems can then be matched to provide additional rejection and higher purity. The L2 farm will consist of around 500 quad-core CPUs; the average processing time at L2 is 40 ms and the expected output rate is around 2 kHz.

The EF is subsequently seeded by L2. The EF has, on average, 4 s to process each event. This allows the use of the more sophisticated offline reconstruction algorithms, as well as offline-like calibration and alignment corrections. The Event Filter algorithms run on a farm of 1800 dual quad-core CPUs processors. The EF an output rate is limited to 200 Hz, assuming an event size

of 1.5 Mb. The execution of the HLT algorithms is organized by the HLT Steering [4] based on the static configuration information and on the dynamic event data. The configuration contains a list of the active signatures (trigger menu) and their thresholds, pass-through fractions and prescale factors. The HLT algorithm sequences are divided into reconstruction steps followed by verification steps. The chain of algorithms can be stopped at any of the verification steps if it is found to be non-viable (early rejection), thus freeing resources for the next signature.

The HLT algorithms are logically divided into groups of related signatures. Currently these are known loosely as: e/γ , μ , τ , jets, b-tag, B physics, missing ET, cosmics and minimum bias. Algorithms from one or more groups, together with configuration information such as threshold values and prescale factors are used to build the trigger signatures that form the trigger menu.

3. Expected Performance in Early Running

During the LHC startup phase, where low luminosity conditions ($10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ or less) and low number of bunches in the machine are expected to prevail, the focus of the trigger selection strategy will be to commission the trigger and the detector and to ensure that established Standard Model processes are observed.

In the very early operations many triggers, especially those at higher thresholds, will operate in pass-through mode, which entails executing the trigger algorithms but accepting the event independent of the decision. This allows the trigger selections and algorithms to be validated to ensure that they are robust against the varying beam and detector conditions that are hard to predict before data taking. As the luminosity increases, the use of higher thresholds, isolation criteria and tighter selections at HLT become necessary to reduce the background rates while achieving selection of interesting physics with high efficiency.

An initial trigger menu has been designed for a luminosity of $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ using simulated minimum-bias events. The following notation is used to label the different trigger items: EM (electromagnetic), J (jets), MU (muons), and tau (tau leptons). Table 1 shows a representative set of L1 signatures, their prescale factors and estimated rates during the start-up for an assumed luminosity of $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$. At low luminosities, a single non-isolated EM trigger with threshold of 7 GeV can be used without the application of prescale factors at the first trigger level with a rate of about 5 kHz. The total L1 output rate out for the EM objects is about 10 kHz. For the jet items, a single trigger with a threshold of 120 GeV can be deployed without prescale factors and has a L1 output rate of about 9 Hz. Single muon and di-muon signatures could all be deployed at $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ without prescale factors with further selection at HLT to control the output rates. Tau triggers at low luminosity are chosen to collect large statistics of W and Z boson decays. The reliance on missing energy end total energy trigger will be small especially during the early running period as it is sensitive to various detector effects that will require time to understand.

Table 1: A representative list of L1 trigger items and estimated rates at $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$

Trigger Item	EM3	EM7	2EM3	MU4	MU20	J10	J120	tau6	tau40
Prescale	60	1	1	1	1	42000	1	750	1
Rate (Hz)	674	4900	6500	1730	20	4	9	19	83

Most of the trigger items at L2 and EF are either executed in pass-through mode or with loose selections. Figure 1 shows a summary of the EF output rates for each trigger group and the cumulative rates, which provide a running total of the rates. The sum of the rates for all the trigger groups is more than the cumulative rates due to overlaps between the groups. The total cumulative rates for this trigger menu, accounting for overlaps between the trigger groups, is 12 kHz at L1, 620 Hz at L2 and 197 Hz at EF. Since there are large uncertainties related to the simulation, the estimated rate out of L1 and L2 is kept well below their respective targets of 40 kHz and 1 kHz available during the LHC startup phase. The EF selections have been tuned to yield the targeted EF output rate compatible with the design output rate of 200 Hz.




Figure 1: HLT unique (black) and cumulative (gray) estimated rates at $10^{31} \text{ cm}^{-2} \text{ s}^{-1}$ for different trigger groups.

4. Commissioning the ATLAS Trigger with Cosmic Rays and First Beam

During the last two years (2007 and 2008), the focus of ATLAS commissioning effort has evolved from single detector operation to combined running and integration. Several combined test (called *Milestone Runs*) with cosmic rays have been scheduled to integrate detector, trigger and data acquisition into one global setup for each group of sub-systems (calorimeters, muon detectors, inner detector and magnets). *Milestone Runs* have been very helpful in the commissioning of the subsystems, in the integration of controls and of the trigger and data acquisition. Moreover, ATLAS was operational when the LHC beams circulated for the first time in September 2008. To cope with different commissioning phases, special trigger menus have been prepared with selection criteria suited to cosmic-ray events, beam-halo and low intensity collisions, and where the trigger is mainly based on L1 items with loose requirements on single tracks and energy clusters. Since the cosmic rays do not come from the interaction point but top of the ATLAS cavern, to increase the acceptance, the trigger condition for the L1 muon trigger was set with the largest possible coincidence windows. This means that the trigger roads for the lower thresholds follow the cabling layout. L1 calorimeter trigger was not designed to be able to trigger on cosmic muons since the noise in each trigger tower is comparable to the signal given by a minimum-ionizing particle. Nonetheless, cosmic rays with energy deposits greater than few GeV could be triggered. Figure 2 shows the integrated cosmic

data recorded by ATLAS with different triggers, versus the time for one of the most active period, i.e. after Sep 13, 2008. For most of the runs the full ATLAS detector (except for CSC) was included in the data acquisition. In some runs the Liquid-Argon Calorimeter was taken out, allowing trigger to accept rates of approximately 500 Hz (corresponding to the high rises in the curves).




Figure 2: Integrated cosmic data rate for ATLAS versus time after Sep 13, 2008.




Figure 3: TGC trigger timing distribution with single-beam with respect to bunch-crossing units (25 ns).

During the cosmics run the HLT infrastructure and algorithms were operational and used in a transparent trigger mode, where events are tagged with HLT results, but not discarded accordingly, as it would happen in standard operation. In addition to this, there is an active HLT selection performed based on tracking algorithms in order to enrich selected streams with tracks useful for timing and alignment studies.

On September 10th 2008, and later on for few days, LHC have successfully circulated the first proton beams with a single bunch. For safety considerations, ATLAS has taken the data with Pixel detector was switched off, and some other subsystems were operated with reduced voltage. The solenoid magnet was off, but the toroid systems were operational.

During this phase, in addition to L1 Calo and Muon, two tools have been set-up for early beams: a beam pick-up from the accelerator (BPTX), and a minimum-bias trigger (MBTS) derived from hodoscopes of plastic scintillators installed on the front face of the end-cap calorimeters. The events recorded by ATLAS with the first beams circulating in LHC can be classified as high-multiplicity events, when the beams were dumped on collimators at 140 m from ATLAS, and halo events, when the collimators were open and fewer particles from beam interactions crossed the detector. These events provide useful checks as, for example, the trigger timing alignment with respect to BPTX. Figure 3 shows the relative time distribution for the different TGC triggers, with the time-of-flight effect particularly visible. The TGC wheels are located at $z = \pm 14$ m, so the two trigger planes generate signals with time difference of about 100 ns, corresponding to 4 bunch crossing units.

5. Conclusions

The three levels of the ATLAS trigger have been designed to handle the LHC rates and occupancies at high luminosity. The ATLAS trigger and its performance have been studied in detail both analyzing the performance with simulated data in the initial LHC scenario and operating the system with cosmic runs. The LHC startup in September 2008 found the ATLAS detector ready; LHC data have been recorded and analyzed providing useful information. ATLAS is looking forward to collect data from LHC collisions.

References

- [1] The ATLAS Collaboration, ATLAS Technical Proposal, CERN/LHCC/94-43, December 1994
- [2] The ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST (2008) 3 S08003;
- [3] ATLAS Collaboration, Level 1 Trigger Technical Design Report, CERN/LHCC/98-14 (1998).
- [4] ATLAS Collaboration, High-Level Trigger, Data Acquisition and Controls Technical Design Report, CERN/LHCC/03-022 (2003).
- [5] R. Spiwojs et al, *The ATLAS level-1 central trigger processor core module (CTP_CORE)*, IEEE Trans. Nucl. Sci. **52** (2005) 3211–3215.